

Letter

# Highest-resolution rotation curve of the inner Milky Way proving the galactic shock wave

Yoshiaki SOFUE\*

Institute of Astronomy, The University of Tokyo, 2-21-1 Mitaka, Tokyo 181-8588, Japan

\*E-mail: [sofue@ioa.s.u-tokyo.ac.jp](mailto:sofue@ioa.s.u-tokyo.ac.jp)

Received 2021 June 11; Accepted 2021 July 11

## Abstract

We present a rotation curve (RC) of the inner Galaxy of the first quadrant at  $10^\circ \leq l \leq 50^\circ$  ( $R = 1.3\text{--}6.2$  kpc) with the highest spatial (2 pc) and velocity ( $1.3 \text{ km s}^{-1}$ ) resolutions. We used  $^{12}\text{CO}(J = 1\text{--}0)$ -line survey data observed with the Nobeyama 45 m telescope at an effective angular resolution of  $20''$  (originally  $15''$ ), and applied the tangent–velocity method to the longitude–velocity diagrams by employing the Gaussian deconvolution of the individual CO-line profiles. A number of RC bumps, or local variation of rotation velocity, with velocity amplitudes  $\sim \pm 9 \text{ km s}^{-1}$  and radial scale length  $\sim 0.5\text{--}1$  kpc are superposed on the mean rotation velocity. The prominent velocity bump and corresponding density variation around  $R \sim 4$  kpc in the tangential direction of the Scutum arm (4 kpc molecular arm) is naturally explained by an ordinary galactic shock wave in a spiral arm with small pitch angle, not necessarily requiring a bar-induced strong shock.

**Key words:** galaxies: individual (Milky Way) — galaxies: rotation curve — ISM: CO line

## 1 Introduction

Various methods to derive the rotation curve (RC), or circular velocities in the Galactic disk, of the Milky Way have been proposed such as the tangent–velocity method (TVM) for a gaseous disk, the radial–velocity plus distance method for stars, the trigonometric method for maser sources, and the disk-thickness method for an HI disk (see reviews by Fich & Tremaine 1991; Sofue & Rubin 2001; Sofue 2017, 2020). A large-scale compilation of RC data has been obtained and is available electronically (Iocco et al. 2015; Huang et al. 2016; Pato & Iocco 2017a, 2017b; Krelowski et al. 2018; Sofue 2020).

The TVM measures the terminal velocity of the gaseous disk of the inner Milky Way inside the Solar circle in the HI and CO line emissions (Burton & Gordon 1978; Clemens 1985; Alvarez et al. 1990; McClure-Griffiths &

Dickey 2007; Marasco et al. 2017). It has the advantage of uniquely determining the galactocentric distance by  $R = R_0 \sin l$  without suffering from uncertainty in the distance measurements, where  $R_0$  is the Solar circle radius. The rotation velocity is given by  $V_{\text{rot}} = v_{\text{LSR}} + V_0 \sin l$ , where  $V_0$  is the Sun’s circular velocity and  $v_{\text{LSR}}$  is the radial velocity of the object with respect to the local standard of rest. In this paper we adopt the Galactic constants of  $(R_0, V_0) = (8.0 \text{ kpc}, 238 \text{ km s}^{-1})$  (Honma et al. 2012, 2015).

A further advantage of using the CO line is that it measures the motion of molecular clouds, which are the most massive and individual objects sharply concentrated near the Galactic plane, having the lowest velocity dispersion among Galactic objects. In other words, CO traces the rotational kinematics of the Galactic disk most precisely with the minimum influence by random motion and velocity dispersion as possessed by other species.

In this paper we apply the TVM to the CO-line data of the inner Galaxy obtained by the FUGIN (Four-beam receiver system Unbiased Galactic-plane Imaging survey with the Nobeyama 45 m telescope; Minamidani et al. 2016; Umemoto et al. 2017).<sup>1</sup> We aim first at providing an RC at the highest spatial (2 pc) and velocity ( $1.3 \text{ km s}^{-1}$ ) resolutions of the inner Milky Way in the first quadrant at  $10^\circ \leq l \leq 50^\circ$ , or at a galactocentric distance of  $R = 1.3$  to 6.1 kpc. It covers the tangential directions of the 4 kpc (Scutum,  $l \sim 30^\circ$ ) arm containing the star-forming complex W 43 and of the 3 kpc expanding (Norma,  $\sim 20^\circ$ ) arm. We then discuss rotational fluctuations in relation to the kinematics of the spiral arms and galactic shock (GS) waves.

## 2 Tangent-velocity method to determine circular velocities


### 2.1 Data

The FUGIN survey covered the Galactic disk in the first quadrant at  $l = 10^\circ \leq 50^\circ$  and  $|b| \leq 1^\circ$ . The  $^{12}\text{CO}(J = 1-0)$  line channel maps had a grid spacing of  $8''.5 \times 8''.5 \times 0.65 \text{ km s}^{-1}$  in the  $(l, b, v_{\text{LSR}})$  space. The effective velocity resolution was  $1.3 \text{ km s}^{-1}$ , the rms noise of the brightness temperature  $T_{\text{B}}$  was  $\sim 1 \text{ K}$ , and the effective angular resolution was  $20''$ , while the original beam of the 45 m telescope at the  $^{12}\text{CO}(J = 1-0)$  frequency was  $15''$ . Although FUGIN data also include  $^{13}\text{CO}$  and  $\text{C}^{18}\text{O}$  lines, we analyze the  $^{12}\text{CO}(J = 1-0)$  line data alone here, because the kinematics of the Galactic molecular disk may not be dependent on the C and O isotopes. Moreover,  $^{13}\text{CO}$  and  $\text{C}^{18}\text{O}$  lines represent higher-density cores, and hence they pick up more patchy structures than those mapped by the  $^{12}\text{CO}(J = 1-0)$  line.

### 2.2 Terminal velocities in the longitude-velocity diagram


Tangent velocities are obtained by tracing the uppermost edges of the emission regions in longitude-velocity (LV) diagrams. Figure 1a shows an LV diagram (brightness temperature  $T_{\text{B}}$  of the  $^{12}\text{CO}(J = 1-0)$  line emission against longitude) in the Galactic plane, and figure 1b shows those around  $l = 31^\circ$  at different latitudes,  $b = 0^\circ$ ,  $0^\circ.5$ , and  $0^\circ.75$ . The apparent upper bounds of the LV ridges, or the terminal velocities, seem to systematically decrease with the latitude; this will be measured more quantitatively in subsection 2.4.

Figure 1c shows an LV diagram across the star-forming complex W 43 in the tangential direction of the 4 kpc arm.



**Fig. 1.** (a) CO-line LV diagrams for  $l = 10^\circ$  to  $50^\circ$  at  $b = 0^\circ$ , smoothed in longitude by a Gaussian beam of  $1'.4$  (10 pixels). Note that  $l$  smoothing is not applied in the analysis for RC. (b) LV diagrams from  $l = 30^\circ$  to  $32^\circ$  at three different latitudes,  $b = 0^\circ$  (top),  $+0^\circ.5$  (middle), and  $+0^\circ.75$ . (c) Enlargement around W 43 in the 4 kpc molecular arm. The terminal edge is inclined by  $\sim +10 \text{ km s}^{-1}$  per degree (red lines) in the opposite sense to that for flat rotation with  $-3.6 \text{ km s}^{-1}$  per degree (dashed line). (Color online)

<sup>1</sup> Five tables of the rotation curve are available in the supplementary data section of the online edition of this article.




**Fig. 2.** Typical FUGIN  $^{12}\text{CO}(J = 1-0)$ -line spectra at  $l = 30:4$  and  $30:8$  (black dots) and their deconvolution into Gaussian components (red lines). The fitted center velocity of the outermost component is defined as the terminal velocity. (Color online)

The intensity distribution is clumpy due to giant molecular clouds and the terminal velocity is locally variable with longitude, exhibiting LV ridges with positive gradients as indicated by the red lines, opposite to the general decrease with longitude shown by the dashed line. The local LV variation will be discussed in detail in subsection 3.4 in relation to the spiral arm and galactic shock wave.

### 2.3 Gaussian deconvolution of line profiles

The simplest way to determine the terminal velocity is to pick up the highest-velocity component after deconvolution of the line profile into many components. Figure 2 shows CO line spectra in the Galactic plane at several longitudes around  $l = 31^\circ$ . Each spectrum can be expressed by the superposition of many components, each represented by a Gaussian profile, as indicated by the red lines. The highest-velocity component is uniquely determined in each profile as the rightmost Gaussian component. We then define the terminal velocity as the center value of the highest-velocity Gaussian component.

Before applying the deconvolution, we smoothed the data cube in the latitude direction by a Gaussian beam of  $\delta l \times \delta b = 25'' \times 60''$  in order to increase the signal-to-noise ratio compared to the original data at  $20'' \times 20''$  resolution without losing much longitudinal or velocity resolution.



**Fig. 3.** (a) Terminal velocities at  $b = 0^\circ$  (black dots),  $0:5$  (red triangles), and  $0:75$  (blue cross) from  $l = 30^\circ$  to  $32^\circ$ . (b) The same, but for  $b = 0^\circ$  with the vertical axis enlarged. (c) Terminal velocities from  $l = 10^\circ$  to  $50^\circ$  at  $b = 0^\circ$ . (Color online)

### 2.4 LV plot of terminal velocities

We apply the Gaussian deconvolution to each spectrum of the  $^{12}\text{CO}(J = 1-0)$  line emission of the CO data cube from the FUGIN survey. Figure 3a shows LV plots where the terminal velocities are plotted against longitude for the longitude range  $30^\circ$ – $32^\circ$ . The bars are the Gaussian-fitted full line widths of the components, which represent intrinsic velocity dispersions of clouds on the order of 5 to  $10 \text{ km s}^{-1}$  observed at the velocity resolution of  $1.3 \text{ km s}^{-1}$ . The LV plots show that the terminal velocities systematically decrease with latitude such that they are highest at  $b = 0^\circ$  (red) and lowest at  $b = 0:75$  in the plot in figure 3. In order to trace maximum terminal velocities in the disk, below we analyze the spectra in the Galactic plane at  $b = 0^\circ$ .

The latitudinal variation of terminal velocity may be attributed to the brightness decreasing with latitude: the

higher the latitude, the higher the possibility of detecting fore-/background emissions as the terminal components due to the decreasing brightness. Alternatively, the decreasing velocity with height from the Galactic plane is real, and represents slower rotation in the upper layer of the molecular disk.

Figure 3b shows the LV plot enlarged in the velocity axis, revealing bumpy variation of the terminal velocity against longitude, with neighboring clumps often exhibiting discrete velocity jumps by 5 to 10 km s<sup>-1</sup>. Such a variation represents proper velocity dispersion among the detected terminal-velocity clouds.

Figure 3c shows the LV plot obtained in this way from  $l = 10^\circ$  to  $50^\circ$ , where the terminal velocities are well determined at  $l \gtrsim 20^\circ$ . Larger scatter at  $l \lesssim 20^\circ$  is not only because of the less sensitive observations, but also due to the intrinsically weak CO-line emission in the innermost region.

## 2.5 FUGIN CO-line rotation curve

Since the terminal velocities read from the data at higher latitudes tend to lead to a lower-velocity rotation curve, we measure here the terminal velocities at  $b \sim 0^\circ$ . Using the LV plot in figure 3c, we thus obtain a plot of maximum terminal velocities against longitude, which we adopt as the circular velocities. Figures 4a and 4b show Gaussian running average plots of the circular velocities, or the rotation curves, where both the radius interval and Gaussian half width were taken to be 2 pc and 100 pc, respectively. In figure 4c we enlarge the velocity axis, and compare the RC with the CO-line  $T_B$ , which is proportional to the local molecular gas density.


During the averaging, we removed data points exceeding  $\pm 40$  km s<sup>-1</sup> from the expected mean value as approximated by a linear function,  $V_{\text{mean}} (\text{km s}^{-1}) = 225 + 11[R (\text{kpc}) - 4] \text{ km s}^{-1}$ , as indicated by the dashed lines, in order to avoid anomalously deviated data.

## 3 Discussion

### 3.1 Comparison with the current RCs

The FUGIN RC coincides well with the current RCs (Sofue 2017, 2020; see the referenced papers therein for RC compilations from the literature) within a few km s<sup>-1</sup> at radii greater than 3.5 kpc. The detailed behavior with bumpy fluctuations is found to be very similar to that of the inner RC of the fourth quadrant from H I tangent-velocity measurements (McClure-Griffiths & Dickey 2007).

However, the FUGIN RC tends to show systematically lower values than the current RCs at radii less than 3.5 kpc.



**Fig. 4.** (a) CO-line RC from  $R = 1.3$  to  $6.2$  kpc after a Gaussian running average with width 2 pc. Scattered data outside the two dashed lines ( $\pm 40$  km s<sup>-1</sup> from the mean) have been removed from the averaging. (b) The same, but with 100 pc width. (c) As (a), but enlarged and compared with  $T_B$  ( $\propto n_{\text{H}_2}$ ) of the  $^{12}\text{CO}(J = 1-0)$  line. (Color online)

The same trend is found in the southern RCs in the H I and CO lines observed in the fourth quadrant (Alvarez et al. 1990; McClure-Griffiths & Dickey 2007). This may be attributed either to a real difference from the current RCs obtained at different resolutions and/or in the different Galactic quadrant, or due to an artifact caused by larger scatter at  $l < \sim 20^\circ$ , where we had less accurate fitting because of the weaker or almost vacant CO emission.

### 3.2 Local RC variations

The RC is superposed by wavy and bumpy fluctuation around the mean with amplitude  $\delta V \sim 10$  km s<sup>-1</sup>. The largest bump is found with its peak at  $R = 4.1$  kpc, associated with the Scutum arm. The velocity increases steeply from 210 to 230 km s<sup>-1</sup> between 3.9 and 4.1 kpc, followed

by a plateau-like enhancement up to  $\sim 5.2$  kpc. The velocity plateau is superposed by two more bumps at  $R = 4.4$  and  $4.9$  kpc. Also, at  $R = 3.1$  kpc there is a velocity depression followed by a jump to the peak at  $3.3$  kpc. These velocity bumps are similar to those found in the fourth quadrant in both their amplitudes and scale lengths (McClure-Griffiths & Dickey 2007).


As well as such prominent bumps, there appear superposed smaller fluctuations of radial scale length  $\delta R \leq \sim 0.1$  pc and velocity amplitudes  $\delta V \sim \pm 2\text{--}3$  km s $^{-1}$ . These small bumps may be attributed to random motions of molecular clouds near the tangent points. As already discussed, the velocity fluctuations found inside  $2.6$  kpc can be attributed to noisy data because of the weak CO-line emission.

### 3.3 Imprint by arms

The RC fluctuations with wave lengths of  $\sim 1\text{--}2$  kpc can be attributed to an imprint by spiral arms associated with non-circular streaming motion (Roberts 1969; Martinez-Medina et al. 2020), or to a local gaseous ring and vacancy (Sofue et al. 2009; McGaugh 2019). In either mechanism, conservation of angular momentum results in deceleration of the rotation velocity for the gas flowing from the inside to the arm, and acceleration for the gas from outside.

Let  $\delta R$  and  $\delta V$  be the radius and velocity displacements from the purely circular motion at  $R$  with  $V_{\text{rot}}$ . Neglecting the second-order quantities, we have  $\delta V/V_{\text{rot}} \simeq -\delta R/R$ . In order to attain the observed velocity acceleration by  $\sim \pm 9$  km s $^{-1}$  for  $V_{\text{rot}} = 220$  km s $^{-1}$ , the gas must be accumulated from a region  $\sim \mp 0.36$  kpc around the density peak in the arm. Such gas flows from both sides of the arm center, causing a steep velocity gradient as observed around  $R \sim 4$  kpc in figure 4 to yield  $dV/dR \sim 70$  km s $^{-1}$  kpc $^{-1}$ . This is observed as the positive gradient of the LV ridge around W 43 in the tangential direction of the 4 kpc (Scutum) arm with  $dv_{\text{LSR}}/dl \sim 10$  km s $^{-1}$  per degree in figure 1c, where the expected gradient of the terminal velocity for flat rotation is negative:  $\sim -3.6$  km s $^{-1}$  per degree.

In figure 4c the brightness temperature  $T_{\text{B}}$  of the  $^{12}\text{CO}(J = 1\text{--}0)$  line at tangential velocities is plotted by the red line. The 4 kpc arm shows up as the sharpest  $T_{\text{B}}$  peak at  $R = 4.0$  kpc, containing the molecular complex and star-forming region W 43 (Kohno et al. 2021). Slightly inside this peak at  $R \sim 3.9$  kpc,  $V_{\text{rot}}$  is observed to attain a local minimum displaced about  $\delta V \sim -10$  km s $^{-1}$  from the mean, indicating deceleration of the gas due to accumulation toward the arm center at the 4 kpc density peak. Opposite behavior with  $\delta V \sim +10$  km s $^{-1}$  at  $4.1$  kpc is observed outside the peak.



**Fig. 5.** (a) Observed (dot with error bar) and theoretical (red line)  $\Delta V$  against radius  $R$ . (b) The same, but for  $T_{\text{B}} (\propto \rho)$ . (c) GS model of  $\delta V$  and  $\rho$  (Roberts 1969) against  $\theta$  (solid) and  $R$  (dashed, schematic). (d) Spiral shocked arms (thick lines) and flow lines (thin lines) representing the Scutum and Sgr arms. The right-hand panel shows the relation between  $\delta R$  and  $\theta$  in the GS model. (Color online)

### 3.4 Galactic shock wave in the Scutum arm

Figure 5a shows the variation of the RC excess from the mean,  $\Delta V = V_{\text{rot}} - V_{\text{mean}}$ , where the mean around the Scutum and Sgr arms was approximated by  $V_{\text{mean}} = 225 + 10[R(\text{kpc}) - 4]$  km s $^{-1}$ . The velocity amplitude is measured to be  $\sim \pm 9$  km s $^{-1}$ . Figure 5b shows the CO brightness temperature  $T_{\text{B}} (\propto n_{\text{H}_2})$ , which is proportional to the local volume density of the molecular gas. The density compression at 4 kpc, which is supposed to be the shock front, is

measured to be  $\rho/\rho_0 \sim T_B/T_{B,\min} \sim 20\text{ K}/2\text{ K} \sim 10$ , and the width is  $\delta R \sim 0.1\text{ kpc}$ .

Figure 5c shows theoretical curves of  $\Delta V$  and  $\rho/\rho_0$  plotted against azimuth angle  $\theta$  along a flow line as reproduced from Roberts (1969). The red dashed lines schematically represent those against the radius. Here, the radial distance  $\delta R$  is related to the azimuth angle  $\theta$  using the plot of a flow line through the spiral arms presented by Roberts (1969). The relation is, here, empirically expressed by  $\delta R \sim \Delta(e^{3x/2} - 1)/(e^{3/2} - 1)$ , where  $x = \theta/\pi$ , and  $\Delta$  is the separation between the arms as explained by figure 5d. Note that the radial variation is much sharper than the azimuthal variation.

The red lines in figures 5a and 5b are calculated radial profiles of the velocity and density based on the galactic shock wave theory, where the shock front is put at  $R = 4\text{ kpc}$  and the arm separation is taken to be  $\Delta = 2\text{ kpc}$ , corresponding to the Scutum and Sgr arms. Here, we recalled the three well-known major arms in the first quadrant, the Norma (3 kpc,  $l \sim 20^\circ$ ), Scutum (4 kpc,  $\sim 30^\circ$ ), and Sgr ( $\sim 50^\circ$ ) arms, with tangential radii  $R \simeq 3, 4$ , and 6 kpc, respectively (Nakanishi & Sofue 2016).

We emphasize that the observed RC and density properties of the Scutum arm are well fitted by the GS model: namely, the velocity bump at 4 kpc followed by a plateau, velocity amplitude  $\pm 9\text{ km s}^{-1}$ , sharp and narrow density peak at 4 kpc, and density compression  $\rho/\rho_0 \sim 10$ . The Scutum arm is, therefore, naturally explained by an ordinary galactic shock wave in a normal spiral arm with potential depth corresponding to  $\sim \pm 9\text{ km s}^{-1}$ .

We comment that this simple view of a spiral arm does not contradict the anticipated bar potential and induced kinematics (Binney et al. 1991; Weinberg 1992; Athanassoula & Bureau 1999), if the bar end is located sufficiently inside  $R \sim 4\text{ kpc}$  and the arm is stretched outside the bar end at a small pitch angle.

## 4 Summary

We analyzed FUGIN CO-line data to determine the rotation curve at the highest resolution ever obtained in the inner Milky Way. The RC is found to be superposed by small-scale variations with velocity gradients as high as several tens of  $\text{km s}^{-1}\text{ kpc}^{-1}$  and amplitude  $\pm 9\text{ km s}^{-1}$ . The local RC bump and density variation toward the 4 kpc arm are

understood as due to a galactic shock wave in a normal spiral arm, not requiring a strong shock by a bar potential.

## Supplementary data

Five tables of the rotation curve data are available at [PASJ](#) online.

## Acknowledgments

Data analysis was carried out at the Astronomy Data Center of the NAOJ. CO data were taken from the FUGIN survey with the Nobeyama 45 m telescope available at (<http://nro-fugin.github.io>).

The author declares, there is no conflict of interest in this work.

## References

- Alvarez, H., May, J., & Bronfman, L. 1990, *ApJ*, 348, 495  
 Athanassoula, E., & Bureau, M. 1999, *ApJ*, 522, 699  
 Binney, J., Gerhard, O. E., Stark, A. A., Bally, J., & Uchida, K. I. 1991, *MNRAS*, 252, 210  
 Burton, W. B., & Gordon, M. A. 1978, *A&A*, 63, 7  
 Clemens, D. P. 1985, *ApJ*, 295, 422  
 Fich, M., & Tremaine, S. 1991, *ARA&A*, 29, 409  
 Honma, M., et al. 2012, *PASJ*, 64, 136  
 Honma, M., Nagayama, T., & Sakai, N. 2015, *PASJ*, 67, 70  
 Huang, Y., et al. 2016, *MNRAS*, 463, 2623  
 Iocco, F., Pato, M., & Bertone, G. 2015, *Nature Phys.*, 11, 245  
 Kohno, M., et al. 2021, *PASJ*, 73, S129  
 Krelowski, J., Galazutdinov, G., & Strobel, A. 2018, *PASP*, 130, 114302  
 McClure-Griffiths, N. M., & Dickey, J. M. 2007, *ApJ*, 671, 427  
 McGaugh, S. S. 2019, *ApJ*, 885, 87  
 Marasco, A., Fraternali, F., van der Hulst, J. M., & Oosterloo, T. 2017, *A&A*, 607, A106  
 Martinez-Medina, L. A., Pichardo, B., & Peimbert, A. 2020, *MNRAS*, 496, 1845  
 Minamidani, T., et al. 2016, *SPIE Proc.*, 9914, 99141Z  
 Nakanishi, H., & Sofue, Y. 2016, *PASJ*, 68, 5  
 Pato, M., & Iocco, F. 2017a, *Astrophys. Source Code Library*, ascl:1711.011  
 Pato, M., & Iocco, F. 2017b, *SoftwareX*, 6, 54  
 Roberts, W. W. 1969, *ApJ*, 158, 123  
 Sofue, Y. 2017, *PASJ*, 69, R1  
 Sofue, Y. 2020, *Galaxies*, 8, 37  
 Sofue, Y., et al. 2019, *PASJ*, 71, S1  
 Sofue, Y., Honma, M., & Omodaka, T. 2009, *PASJ*, 61, 227  
 Sofue, Y., & Rubin, V. 2001, *ARA&A*, 39, 137  
 Umemoto, T., et al. 2017, *PASJ*, 69, 78  
 Weinberg, M. D. 1992, *ApJ*, 384, 81