

ISO DEEP FAR-INFRARED SURVEY IN THE LOCKMAN HOLE A SEARCH FOR OBSCURED OBJECTS AT HIGH REDSHIFT

K. Kawara¹, Y. Sato^{2,3}, H. Matsuhara², Y. Taniguchi⁴, H. Okuda², Y. Sofue¹
T. Matsumoto², K. Wakamatsu⁵, L.L. Cowie⁶, R.D. Joseph⁶, & D.B. Sanders⁶

¹Institute of Astronomy, The University of Tokyo, 2-21-1 Osawa Mitaka, Tokyo, 181, Japan.

²Institute of Space and Astronautical Science (ISAS), 3-1-1 Yoshinodai, Sagamihara, Kanagawa, 229, Japan.

³ISO Data Centre, ESA Astrophysics Division, Villafranca del Castillo, Spain.

⁴Astronomical Institute, Tohoku University, Aoba, Sendai 980-77, Japan.

⁵Department of Physics, Gifu University, Gifu 501-11, Japan.

⁶Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA.

ABSTRACT

Two $44' \times 44'$ fields in the Lockman Hole were mapped at 95 and 175 μm using ISOPHOT. A simple program code combined with PIA works well to correct for the drift in the detector responsivity. The number density of 175 μm sources is 3 - 10 times higher than expected from the no-evolution model. The source counts at 95 and 175 μm are consistent with the cosmic infrared background.

Key words: Galaxies; evolution, Galaxies; formation, quasars; general, Galaxies; starburst, Cosmology; observations, Infrared; galaxies.

The combinations of the filters and the detector arrays which were used for the observations are given in Kawara et al. (1998).

Raster mapping was performed with the AOT PHT22 staring raster map mode using 35.4 hours, which were executed between revolutions 194 and 215 (May 28 and June 19, 1996). Each of the two fields is made up of four sub-fields having an approximately $22' \times 22'$ area each. The parameters used for raster mapping of a single sub-field and details of the read-out settings are given in Kawara et al. (1998).

3. DATA REDUCTION

1. INTRODUCTION

As part of the Japan/UH cosmology program using the ISAS guaranteed time, 95 and 175 μm surveys were conducted in the Lockman Hole to search for obscured infrared galaxies at high redshift. The first results and preliminary discussion were presented by Kawara et al. (1998). This paper presents observations, image processing, and discussion on the nature of far-infrared sources. Point source extraction from the maps and source counts will be presented in a forthcoming paper (Kawara et al. 1999).

2. OBSERVATIONS

We have carried out a deep far-infrared survey at 95 and 175 μm by using ISOPHOT (Lemke et al. 1996) which is an imaging photopolarimeter on board the Infrared Space Observatory (ISO; Kessler et al. 1996). Two fields named LHEX and LHNW, each extending approximately $44' \times 44'$, were selected in the "Lockman Hole", a region with the smallest HI column density in the sky (Lockman et al. 1986), to minimize confusion noise due to infrared cirrus.

The PHT Interactive Analysis (PIA) - Gabriel et al. (1997) - version 6.4 was used with the default settings to reduce the edited raw data created in November, 1996 via the off-line processing version 5.1 or 5.2. Maps of 95 μm created from the PIA's Astronomical Analysis Processing (AAP) are dominated by noise due to drift in the detector responsivity. The AAP created maps are thus cleaned by a median smooth routine developed by Kawara et al. (1998).¹ The photometric error introduced from this correction is 4% or better and no extended brightness structures recognized within a $22' \times 22'$ area are affected.

The flux calibration can be made using either the COBE background measurements or the IRAS point source fluxes. Note that the intensity of interplanetary dust emission varies little at such high ecliptic latitude ($\beta \sim 45^\circ$) as the solar elongation changes. The fields contain only one IRAS source (F10507+5723). This is the brightest source detected in our survey and identified with a Sb galaxy UGC 06009 (Thuan & Sauvage 1992). The IRAS fluxes are 533 mJy at 60 μm and 1218 mJy at 100 μm (IRAS FSC). Its flux ratio $f(100 \mu\text{m})/f(60 \mu\text{m}) =$

¹This routine is now included in PIA and called "Median Filtering."

2.29 is fitted to a combination of the infrared cirrus and starburst spectra given by Pearson & Rowan-Robinson (1996), if 76% of the $100\ \mu\text{m}$ flux comes from the cirrus component. This predicts $f(175\ \mu\text{m}) = 1133\ \text{mJy}$ from $f(175\ \mu\text{m})/f(100\ \mu\text{m}) = 0.93$ which is comparable to that in the inner arm of a Sc galaxy M101 (Hippelein et al. 1996). The COBE based scaling yields $347\ \text{mJy}$ at $95\ \mu\text{m}$ and $451\ \text{mJy}$ at $175\ \mu\text{m}$, implying that the COBE based scaling may underestimate point source fluxes by 3.5 at $95\ \mu\text{m}$ and 2.5 at $175\ \mu\text{m}$. The origin of these differences is unclear. In the following discussion, the ISO fluxes are scaled based on the IRAS measurements.

4. DISCUSSION

There are numerous spots in the 95 and $175\ \mu\text{m}$ maps presented by Kawara et al. (1998). Bright spots cannot be attributed to peaks of the infrared cirrus. The total HI column density N_H in our fields is approximately $6 \times 10^{19}\ \text{cm}^{-2}$ (Jahoda et al. 1990). The correlation between N_H and the infrared cirrus brightness indicates $B_{\text{cirrus}}(95\ \mu\text{m}) = 0.41\ \text{MJy sr}^{-1}$ and $B_{\text{cirrus}}(175\ \mu\text{m}) = 0.81\ \text{MJy sr}^{-1}$ (Table 2 in Boulanger et al. 1996). Using the models by Gautier et al. (1992) with $\alpha = -2.9$ for the index of the spatial power spectrum of the infrared cirrus, $\sigma_{\text{cirrus}}(95\ \mu\text{m}) \sim 0.14\ \text{mJy}$ and $\sigma_{\text{cirrus}}(175\ \mu\text{m}) \sim 1.7\ \text{mJy}$ are obtained for one sigma cirrus confusion noise. The $3\ \sigma$ flux levels reached in our survey are tentatively estimated to be $\sim 45 \pm 20\ \text{mJy}$ at $95\ \mu\text{m}$ and $\sim 45 \pm 25\ \text{mJy}$ at $175\ \mu\text{m}$. Hence, a probability of having a cirrus peak brighter than the detection limits is so low that most of spots are likely to be galaxies.

In the two fields LHEX and LHNW, covering 1.1 square degrees in total, there are 36 and 45 sources brighter than $150\ \text{mJy}$ at 95 and $175\ \mu\text{m}$, respectively. Out of 45 $175\ \mu\text{m}$ sources, 36 sources are within the area which was observed at both wavelengths. 31 of the 36 sources have the $95\ \mu\text{m}$ counterparts within $40''$ that were detected above the $3\ \sigma$ flux level. The completeness down to $150\ \text{mJy}$ is estimated to be almost 100%, because of the low probability of cirrus confusion and the high SNR (signal to noise ratios) detection (typical SNR ~ 10). The cumulative counts down to $150\ \text{mJy}$ are thus $1.1 \times 10^5\ \text{sources sr}^{-1}$ at $95\ \mu\text{m}$ and $1.3 \times 10^5\ \text{sources sr}^{-1}$ at $175\ \mu\text{m}$. Fig. 1 compares the 95 and $175\ \mu\text{m}$ counts with the models developed by Guiderdoni et al. (1997, 1998). The number density of $175\ \mu\text{m}$ sources brighter than $150\ \text{mJy}$ are ten times higher than that expected from the no-evolution model. The counts both at 95 and $175\ \mu\text{m}$ counts are consistent with the models containing a strongly evolving component which is required to account for the cosmic infrared background found by Puget et al. (1996).

ACKNOWLEDGEMENTS

The ISOPHOT data presented in this paper was reduced using PIA, which is a joint development by the ESA Astrophysics Division and the ISOPHOT con-




Figure 1. The cumulative number counts at 95 and $175\ \mu\text{m}$ down to $150\ \text{mJy}$ which are compared with the models by Guiderdoni et al. The no-evolution model is plotted with the dash-dot line, and scenarios E and A with the solid and dash lines, respectively.

sortium. The ISOPHT consortium is led by the Max Planck Institute for Astronomy (MPIA), Heidelberg.

REFERENCES

- Boulanger, F., Abergel, A., Bernard, J.-P., et al. 1996, A&A 312, 256.
- Gabriel, C., et al. 1997, Proc. of the ADASS VI Conference, G. Hunt & H.E. Payne (eds), pp. 108
- Gautier, T.N., Boulanger, F., P. Pérault, M., & Puget, J.L. 1992, AJ, 103, 1313
- Guiderdoni, B., Bouchet, F.R., Puget, J.-L., Guilaïne, G., & Hivon, E. Nature, 390, 257
- Guiderdoni, B., Hivon, E., Bouchet, F.R., & Maffei, B. MNRAS, 295, 877
- Hippelein, H., Lemke, D., Haas, M., et al. 1996, A&A, 315, L82
- IRAS FSC, version 2.0, 1990, Joint IRAS Science Working Group, U.S. Government Printing Office, Washington D.C.
- Jahoda, K., Lockman, F.J., & McCammon, D. 1990, ApJ, 354, 184
- Kawara, K., Sato, Y., Matsuhara, H. et al. 1998, A&A 336, L9
- Kawara, K. et al. 1999, in preparation.
- Kessler, M.F., Steinz, J.A., Anderegg, M.E. et al., 1996, A&A 315, L27
- Lemke, D., Klaas, U., Abolins, J., et al. 1996, A&A 315, L64
- Lockman, F.J., Jahoda, K., & McCammon, D. 1986, ApJ, 302, 432
- Pearson, C. & Rowan-Robinson, M. 1996, MNRAS, 283, 174
- Puget, J.L. et al. 1996, A&A, 308, L5
- Thuan, T.X., & Sauvage, M. 1992, A&AS, 92, 749